

Even More Plural Theory in HOL

Carl Pollard

September 13 & 20, 2018

Game Plan

- At first, we work in HOL with basic types e and t .
- For A a type, an ‘ A -set’ means something of type $A \rightarrow t$.
- We introduce a Link-ish theory, using a unary type constructor **Agg** of **aggregates**.
- We make **Agg** into a monad.
- We then elaborate the theory to classify predicates and handle ‘fancy’ plurals.
- Eventually the theory will have to be framed in a richer type theory (at least with dependent sums indexed by the natural numbers) in order to handle predicates that can be predicated *only* of plurals.
- (Hyper-)intensionality will have to wait.

Review of Useful Defined Terms for Set-ish Business

$$\{-\}_A := \lambda xy : A. x = y$$

$$\text{nonempty}_A := \lambda S : A \rightarrow \mathbf{t}. \exists x : A. S x$$

$$\text{singleton}_A := \lambda S : A \rightarrow \mathbf{t}. \exists x : A. S = \{x\}$$

$$\text{pluralton}_A := \lambda S : A \rightarrow \mathbf{t}. \exists xy : A. (x \neq y) \wedge (S x) \wedge (S y)$$

$$\text{injective}_{A,B} := \lambda f : A \rightarrow B. \forall xy : A. [(f x) = (f y)] \rightarrow x = y$$

$$\subseteq_A := \lambda ST : A \rightarrow \mathbf{t}. \forall x : A. (S x) \rightarrow (T x)$$

$$\bigcup_A := \lambda S : (A \rightarrow \mathbf{t}) \rightarrow \mathbf{t}. \lambda x : A. \exists T : A \rightarrow \mathbf{t}. (S T) \wedge (T x)$$

$$\bigcup_A := \lambda ST : A \rightarrow \mathbf{t}. \lambda x : A. (S x) \vee (T x)$$

Link-isch Theory Basics (1/2)

- We introduce a unary type constructor \mathbf{Agg} of *aggregates*.
- We introduce the type-indexed family of constants

$$\mathbf{atoms}_A : (\mathbf{Agg} A) \rightarrow A \rightarrow \mathbf{t}$$

axiomatized as bijections from the A -aggregates to the nonempty A -sets:

$$\vdash \text{injective } \mathbf{atoms}_A$$

$$\vdash \forall m : \mathbf{Agg} A. \text{nonempty } (\mathbf{atoms} m)$$

$$\vdash \forall S : A \rightarrow \mathbf{t}. (\text{nonempty } S) \rightarrow \exists m : \mathbf{Agg} A. S = (\mathbf{atoms} m)$$

Link-isch Theory Basics (2/2)

- We define our counterpart of Link's part-of order as follows:

$$\sqsubseteq_A := \lambda mn : \mathbf{Agg} \ A. (\text{atoms } m) \subseteq (\text{atoms } n)$$

which makes the bijection from A -aggregates to nonempty A -sets into an order-isomorphism.

- We define singular and plural aggregates straightforwardly:

$\text{singular}_A := \lambda m : \mathbf{Agg} \ A. \text{singleton } (\text{atoms } m)$

$\text{plural}_A := \lambda m : \mathbf{Agg} \ A. \text{pluralton } (\text{atoms } m)$

The Aggregate Monad

The nonempty powerset functor has a well-known monad structure (aka the nondeterminism monad), which we transfer to \mathbf{Agg} via the atoms bijection:

Unit:

$$\eta_A : A \rightarrow (\mathbf{Agg} A)$$
$$\vdash \forall x : A. (\text{atoms } (\eta_A x)) = \{x\}$$

Multiplication:

$$\mu_A : (\mathbf{Agg}^2 A) \rightarrow (\mathbf{Agg} A)$$
$$\vdash \forall m : \mathbf{Agg}^2 A. \text{atoms } (\mu_A m) =$$
$$\bigcup (\lambda S : A \rightarrow \mathbf{t}. \exists n : \mathbf{Agg} A. (\text{atoms } m n) \wedge (S = (\text{atoms } n)))$$

Functor at level of terms:

$$\text{agg}_{A,B} : (A \rightarrow B) \rightarrow (\mathbf{Agg} A) \rightarrow (\mathbf{Agg} B)$$
$$\vdash \forall f : A \rightarrow B. \forall m : \mathbf{Agg} A. \text{atoms } (\text{agg}_{A,B} f m) =$$
$$\lambda y : B. \exists x : A. (\text{atoms } m x) \wedge y = (f x)$$

Aggregate Sum

- We introduce a family of constants corresponding to Link's (binary) sum:

$$\vdash \sqcup_A : (\mathbf{Agg} A) \rightarrow (\mathbf{Agg} A) \rightarrow (\mathbf{Agg} A)$$

$$\vdash \forall mn : \mathbf{Agg} A. (\mathbf{atoms} (m \sqcup n)) = (\mathbf{atoms} m) \cup (\mathbf{atoms} n)$$

- The new axiom schema makes the order isomorphisms from aggregates to nonempty sets of atoms into join-semilattice isomorphisms.
- We lack a counterpart to Link's infinitary sum (so the join semilattices of aggregates are not complete).
- We didn't really need infinitary sums anyway.

Nonquantificational NPs

- As in traditional accounts, we translate names of entities with constants of type e , e.g. $j : e$ (*John*), $m : e$ (*Mary*).
- *And* is treated as ambiguous between its familiar boolean meaning (for conjoining truth values or functions with final result type t) and the new meaning \sqcup .
- Entities can't be summed, but the corresponding singular aggregates can, e.g. $(\eta j) \sqcup (\eta m) : \text{Agg } e$ (*John and Mary*).

Indifferent Predicates

- Predicates which can predicate of both singulars and plurals, such as *performed*, are treated as sets of aggregates, i.e. (for entities) $(\text{Agg } e) \rightarrow t$:

perform $((\eta m) \sqcup (\eta j))$ (*Mary and John performed.* [as a unit])

perform (ηm) (*Mary performed.*)

- But *Mary and John performed* also has a distributive) reading, usually expressed using boolean conjunction. We'll come back to that.
- And *Mary performed* is standardly analyzed as having an entity predicate (type $e \rightarrow t$). We'll come back to that too.

Distributivity (1/2)

- We define an aggregate predicate to be *distributive* provided it holds of an aggregate iff it holds of all the aggregate's singular subparts:

$$\text{distrib}_A := \lambda T : (\text{Agg } A) \rightarrow \mathbf{t} . \forall m : \text{Agg } A . (T \ m) \leftrightarrow (\forall n : \text{Agg } A . ((\text{singular } n) \wedge (n \sqsubseteq m)) \rightarrow (T \ n))$$

- We analyze distributive predicates (e.g. *die*) as aggregate predicates which are *axiomatically* distributive:

$$\begin{aligned} \vdash \text{die} : (\text{Agg } e) \rightarrow \mathbf{t} \\ \vdash (\text{distrib } \text{die}) \end{aligned}$$

Distributivity (2/2)

- For each type A we can define two functions that set up a bijection between the A -predicates and the distributive $(\text{Agg } A)$ -predicates, called **individualization** and **distributivization**:

$$\text{indiv}_A := \lambda T : (\text{Agg } A) \rightarrow \text{t}.\lambda x : A.T(\eta x)$$

$$\text{dist}_A := \lambda S : A \rightarrow \text{t}.\lambda m : \text{Agg } A.\forall x : A.(\text{atoms } m x) \rightarrow (S x)$$

Indifferent Predicates Revisited

- Any aggregate predicate T can be mapped to a distributive predicate, namely $\text{dist}(\text{indiv } T)$.
- For example, the distributive reading of *Mary and John performed* can be expressed (*without* boolean conjunction):

$$\text{dist}(\text{indiv perform}) ((\eta m) \sqcup (\eta j))$$

- Also, *Mary performed* can be expressed with an entity predicate:

$$\text{indiv perform } m$$

Singular and Plural Nouns (1/4)

N.B.: Here, by ‘noun’, we really mean ‘count noun’.

- On a first pass, we’ll treat (entity-)plural noun denotations as distributive aggregate predicates $((\text{Agg } e) \rightarrow t)$ and singular nouns as their individualizations (*a fortiori*, entity predicates $(e \rightarrow t)$):

$$\vdash \text{bees} : (\text{Agg } e) \rightarrow t$$
$$\vdash \text{distrib bees}$$
$$\text{bee} := (\text{indiv bees}) : e \rightarrow t$$

- $\text{bees } ((\eta e) \sqcup (\eta d))$ (*Eric and Derek are bees.*)
 $\text{bee } s$ (*Sam is a bee.*)

Singular and Plural Nouns (2/4)

- For some common nouns such as *swarm*, the *singular* form already denotes a predicate of aggregates, which moreover holds only of plurals. We analyze the corresponding plural nouns as denoting aggregates of aggregates:

$$\vdash \text{swarms} : (\text{Agg}^2 e) \rightarrow t$$

$$\vdash \text{distrib swarms}$$

$$\text{swarm} := (\text{indiv swarms}) : (\text{Agg } e) \rightarrow t$$

$$\vdash \forall m : \text{Agg } e. (\text{swarm } m) \rightarrow (\text{plural } m)$$

- $\text{swarm } ((\eta e) \sqcup (\eta d) \sqcup (\eta b) \sqcup (\eta s))$ (*Eric, Derek, Buzz, and Sam are a swarm.*)
 $\text{swarm } (\eta e)$ (*Eric is a swarm.*)
(merely false; cf. * *Eric is bees*)

Singular and Plural Nouns (3/4)

- This treatment of plural nouns isn't quite right, because entity-plural noun denotations can't hold of entities, or even of singular aggregates:
 - a. Eric/the bee is/are/wants to be bees.
- Rather, they (and nondistributive plural predicates, such as *be alike* and *hate each other*) can only hold of *plural* aggregates (*John and Mary, the children*).
- In fact, it seems we really should say something stronger: that they can only be *predicated* of plural aggregates.
- But as yet we can't formalize that idea, because there are no *types* of plural aggregates.

Singular and Plural Nouns (4/4)

- The following examples aren't merely false:
 - a. The honeybee/Eric is/are/wants to be bumblebees.
 - b. The farmer/Pedro is/are alike.
 - c. The mathematician/Fermat hated each other.
- Negating them does not improve them.
- We'll ignore this issue for now; we'll eventually resolve it by adding a separate type constructor for plurals .
- But not today.

Definites (1/2)

- We assume *the* is ambiguous:

$\text{the}_A^{\text{sng}} : (A \rightarrow t) \rightarrow A$, which presupposes a contextually salient member of the argument predicate and returns it.

$\text{the}_A^{\text{plu}} : ((\text{Agg } A) \rightarrow t) \rightarrow (\text{Agg } A)$, which presupposes a contextually salient *plural* member of the predicate and returns it.

- Note that $\text{the}_{(\text{Agg } A)}^{\text{sng}}$ and $\text{the}_A^{\text{plu}}$ have the same type but different presuppositions.
- [Eric, Derek, Buzz, and Sam]₁ were bees. They₁ gave a four-hour joint presentation on waggle dance semantics. Then [the exhausted swarm]₁ returned to it₁s colony.

Definites (2/2)

- $(\text{the}_e^{\text{sng}} \text{ bee}) : e$
 $(\text{the}_e^{\text{plu}} \text{ bees}) : \text{Agg } e$
 $(\text{the}_{(\text{Agg } e)}^{\text{sng}} \text{ swarm}) : \text{Agg } e$
 $(\text{the}_{(\text{Agg } e)}^{\text{plu}} \text{ swarms}) : \text{Agg}^2 e$
- $(\text{the}_e^{\text{plu}} \text{ bees}) \sqcup_e (\text{the}_e^{\text{plu}} \text{ wasps}) : \text{Agg } e$ (an aggregate each of whose atoms is either one of the bees or one of the wasps)
 $(\eta_{(\text{Agg } e)} (\text{the}_e^{\text{plu}} \text{ bees})) \sqcup_{(\text{Agg } e)} (\eta_{(\text{Agg } e)} (\text{the}_e^{\text{plu}} \text{ wasps})) : \text{Agg}^2 e$
(an aggregate with two atoms: the bees and the wasps)

Nondistributable Plural Predicates (1/3)

- *Nondistributable* plural predicates differ from plural common nouns in having no individual counterparts:
 - a. The bees/Sam and Buzz are alike/converged/buzzed each other.
 - b. *The bee/Sam is alike/converged/buzzed each other.
- Some nondistributable plural predicates aren't fussy about what their arguments are plurals *of*:
 - c. Eric and Derek/juggling and miming/donkeys and burros/the Riemann Hypothesis and the Goldbach Conjecture/17 and 37/conjunction and sum are alike.
- We can analyze such predicates as families of type-indexed (ordinary) predicates, e.g.

$$\text{alike}_A, \text{converge}_A : (\text{Agg } A) \rightarrow t$$

Nondistributable Plural Predicates (2/3)

- $\text{converge}_e : (\text{Agg } e) \rightarrow t$
 $\text{converge}_{(\text{Agg } e)} : (\text{Agg}^2 e) \rightarrow t$
 $(\text{dist } \text{converge}_e) : (\text{Agg}^2 e) \rightarrow t$
- $(\text{converge}_e ((\eta s) \sqcup (\eta b)))$ (*Sam and Buzz converged.*)
 $(\text{converge}_e (\text{the}_{(\text{Agg } e)}^{\text{sg}} \text{swarm}))$ (*The swarm converged.*)
- $(\text{converge}_{(\text{Agg } e)} (\text{the}_{(\text{Agg } e)}^{\text{plu}} \text{swarms}))$ (*The swarms converged.*) [They all headed to the same location.]
 $(\text{dist } \text{converge}_e (\text{the}_{(\text{Agg } e)}^{\text{plu}} \text{swarms}))$ (*The swarms converged.*) [Each of them converged.]
 $(\text{converge}_e (\mu_e (\text{the}_{(\text{Agg } e)}^{\text{plu}} \text{swarms})))$ (*The swarms converged.*) [The bees in the swarms all headed to the same location.]

Nondistributable Plural Predicates (3/3)

- $\text{alike}_e : (\text{Agg } e) \rightarrow t$
 $\text{alike}_{(\text{Agg } e)} : (\text{Agg}^2 e) \rightarrow t$
 $(\text{dist } \text{alike}_e) : (\text{Agg}^2 e) \rightarrow t$
- $(\text{alike}_e ((\eta s) \sqcup (\eta b)))$ (*Sam and Buzz are alike.*)
 $(\text{alike}_e (\text{the}_e^{\text{plu}} \text{ bees}))$ (*The bees are alike.*)
- Abbreviations:
 $\text{bw} := (\text{the}_e^{\text{plu}} \text{ bees}) \sqcup_e (\text{the}_e^{\text{plu}} \text{ wasps})$
 $\text{BW} := (\eta_{(\text{Agg } e)} (\text{the}_e^{\text{plu}} \text{ bees})) \sqcup_{(\text{Agg } e)} (\eta_{(\text{Agg } e)} (\text{the}_e^{\text{plu}} \text{ wasps}))$
- $(\text{alike}_{(\text{Agg } e)} \text{ BW})$ (*The bees and the wasps are alike.*) [They are similar aggregates.]
 $(\text{dist } \text{alike}_e \text{ BW})$ (*The bees and the wasps are alike.*) [The bees are alike, and so are the wasps.]
 $(\text{alike}_e \text{ bw})$ (*The bees and the wasps are alike.*) [The insects, which comprise the bees and the wasps, are alike.]